

Impact of Structured Massage Therapy on Physiological Stability, Comfort, Sleep, and Stress in Late Preterm Infants: A Randomized Controlled Trial

Rıdvan Akdoğan¹, Yusuf Gürgan¹, Mehmet Emin Düken²

¹Department of Pediatric Nursing, Faculty of Health Sciences, Van Yüzyüncü Yıl University, Van, Türkiye; ²Department of Child Health and Diseases Nursing, Health Sciences of Faculty, Harran University, Şanlıurfa, Türkiye

Abstract:

Objective: This randomized controlled trial aimed to evaluate the effects of structured massage therapy on sleep quality, stress, comfort, and physiological parameters in late preterm infants.

Methods: Forty late preterm infants (35–37 gestational weeks) were randomly allocated to a massage group (n=20) or a control group (n=20). The intervention group received a standardized 15-minute massage three times daily for three consecutive days. Data were collected using physiological monitoring, the Premature Infant Comfort Scale (PICS), the Neonatal Stress Scale (NSS), and actigraphy (SOMNOwatch™). This study was conducted in the neonatal intensive care units of Van Training and Research Hospital, University of Health Sciences, between September 2024 and September 2025, and was approved by the relevant institutional ethics committee in accordance with the Declaration of Helsinki.

Results: Massage therapy significantly improved physiological stability, demonstrated by lower heart rate and respiratory rate and higher oxygen saturation compared to controls ($P<0.05$). Post-intervention comfort scores were significantly higher in the massage group ($P=0.002$). Significant within-group improvements were observed in sleep and stress parameters ($P<0.001$); however, no statistically significant between-group differences were identified ($P>0.05$).

Conclusion: Massage therapy is an effective and clinically relevant non-pharmacological intervention for enhancing physiological stability and comfort in late preterm infants. Although improvements in sleep and stress were comparable to routine care, the significant improvement in comfort supports its integration into neonatal nursing protocols to optimize immediate clinical outcomes and neurobehavioral organization.

Clinical Trials Number: NCT07322731

Keywords: Massage Therapy, Preterm Infant, Health Care, Physiological Stability

Approximately 15 million infants are born preterm (gestational age <37 weeks) worldwide each year, a figure that continues to rise as a global public health concern [1]. Complications of preterm birth are responsible for 35% of neonatal deaths and carry significant long-term morbidity risks among survivors, including neurodevelopmental disorders, learning disabilities,

Submitted: March 14, 2026 Accepted: May 24, 2026 Published Online: June 17, 2026

How to cite this article: Akdoğan R, Gürgan Y, Düken ME. Impact of Structured Massage Therapy on Physiological Stability, Comfort, Sleep, and Stress in Late Preterm Infants: A Randomized Controlled Trial. Eur Res J. 2026. doi: 10.18621/eurj.1301

Corresponding author: Rıdvan Akdoğan, PhD., Assist. Prof., Phone: +90 432 225 11 80, E-mail: ridvanakdogan@yyu.edu.tr

This is an open-access article distributed under the terms of a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits any non-commercial use, sharing, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if you modified the licensed material. You do not have permission under this licence to share adapted material derived from this article or parts of it.

Available Online at <https://www.eurj.org.tr>



and visual and auditory impairments [1, 2]. This vulnerable population requires prolonged monitoring in the Neonatal Intensive Care Unit (NICU) to sustain vital functions such as feeding and respiration [3].

The NICU environment constitutes an overstimulating and stressful setting for the immature nervous systems of preterm infants. These infants are continuously exposed to multiple painful and distressing stimuli, including elevated noise levels, persistent light exposure, and invasive medical procedures [4]. Such environmental factors not only destabilize vital signs — including heart rate, respiratory rate, and oxygen saturation — but may also inflict lasting damage on the autonomic nervous system [5, 6]. Furthermore, intensive care routines in the NICU significantly disrupt sleep cycles, which are critical for brain development and synaptic plasticity [7, 8]. Sleep disruption can adversely affect growth hormone secretion, thereby delaying clinical recovery and prolonging the time to hospital discharge [7].

In contemporary neonatal nursing practice, non-pharmacological approaches have come to the forefront as strategies for enhancing infant comfort, managing stress, and promoting stabilization. Alongside methods such as kangaroo care, environmental modifications, and olfactory stimulation, infant massage has attracted growing attention as a safe and cost-effective complementary intervention [9, 10]. The literature indicates that structured massage protocols augment vagal activity, thereby reducing cortisol levels and contributing to stress reduction and improved sleep quality [11, 12]. Meta-analyses and randomized controlled trials have further demonstrated that pediatric massage prolongs sleep duration [13], supports weight gain [14], and shortens the length of hospital stay [15].

Nevertheless, evidence-based studies that simultaneously examine the synergistic effects of infant massage on stress, comfort, and vital signs in preterm infants — particularly within the late preterm cohort (35–37 weeks) — remain limited. It is of paramount importance that NICU nurses integrate non-pharmacological approaches of this nature into clinical practice in accordance with the principles of atraumatic care, rather than focusing exclusively on medical management [16]. In this context, the present randomized controlled trial aimed to comprehensively evaluate the effects of massage therapy on sleep

quality, comfort, stress indicators, and physiological parameters in infants born at 35–37 weeks of gestation. The following hypotheses were tested: massage therapy in preterm infants would (1) support physiological stabilization, (2) reduce stress and enhance comfort, and (3) improve sleep architecture (duration and efficiency).

H1: Post-intervention heart rate, respiratory rate, and oxygen saturation (SpO₂) values in massaged preterm infants will be statistically significantly more stable than those in the control group.

H2: Massage therapy will statistically significantly improve comfort levels (Premature Infant Comfort Scale [PICS] scores) in preterm infants compared to the control group.

H3: Stress scores (Neonatal Stress Scale [NSS]) in massaged preterm infants will be statistically significantly lower than those in the control group.

H4: Massage therapy will significantly increase sleep efficiency and total sleep duration while reducing the number of awakenings and wake after sleep onset (WASO) duration compared to the control group.

METHODS

Study Design and Aim

This study was conducted as a randomized controlled experimental trial to evaluate the effects of massage therapy on sleep quality, comfort, stress indicators, and physiological parameters in late preterm infants (35–37 gestational weeks) admitted to the NICU. The study was designed and reported in accordance with the CONSORT (Consolidated Standards of Reporting Trials) guidelines (Figure 1).

Sample and Power Analysis

The study population consisted of preterm infants admitted to the Level I and Level II NICU services of Van Training and Research Hospital, University of Health Sciences, between September 2024 and September 2025. Sample size was calculated using G*Power (v3.1.9.2), based on the reference study by Düken and Yayan [17] conducted with preterm infants. Using an effect size of $d=1.192$, $\alpha=0.05$, and 95% power ($1-\beta$), the analysis determined that a minimum




FIGURE 1. CONSORT flow diagram [36].

of 20 infants per group were required. The study was completed with a total of 40 infants (20 in the massage group, 20 in the control group). Of the 54 infants assessed for eligibility, 14 were excluded (8 did not meet the inclusion criteria and 6 declined participation), resulting in 40 infants being randomized into the study groups (n=20 massage, n=20 control).

Randomization and Blinding

Simple randomization — a method equivalent to complete randomization in its balancing properties — was employed for group allocation. For each eligible infant, the researcher drew a ball from a sealed black pouch using a closed-selection method. White balls designated the massage group and red balls designated the control group, thereby minimizing selection bias

and ensuring homogeneous distribution between groups. No dropouts occurred during the study, and all randomized participants were included in the analyses.

Given the nature of the intervention, blinding of the interventionist was not feasible. However, assessor blinding was implemented to reduce measurement bias. The researcher responsible for statistical analyses received data coded as "Group A" and "Group B," with group identities remaining concealed until all analyses were complete.

Inclusion and Exclusion Criteria

Inclusion Criteria

Parental written informed consent; gestational age of 35–37 weeks (up to 36 weeks + 6 days); birth weight >1,500 g; Apgar score ≥ 7 ; absence of known congenital or neurological disorders.

Exclusion Criteria

Requirement for mechanical ventilatory support; history of major surgery; active infection; or any dermatological condition compromising skin integrity that would preclude massage application.

Data Collection Instruments

Mother and Infant Demographic Information Form

A structured form capturing demographic, obstetric, and neonatal characteristics (gestational age, birth weight, sex, etc.).

Vital Signs Recording Form

This form was developed by the researcher based on the relevant literature to assess the physiological status of preterm infants [18,19]. It includes key vital parameters: heart rate, respiratory rate, body temperature, and oxygen saturation. Measurements were obtained using standard clinical monitoring equipment and recorded at predetermined time points.

Premature Infant Comfort Scale (PICS)

The PICS is an adapted form of the COMFORT scale developed to assess comfort levels in preterm infants [20, 21]. Its Turkish validity and reliability were established by Alemdar and Tüfekçi [22]. The scale comprises 7 parameters — alertness, agitation, respiration/crying, body movement, muscle tone, facial expression, and heart rate — each scored from 1 to 5. Total scores range from 7 to 35, with higher scores indicating lower comfort. The cutoff score is ≥ 17 .

Neonatal Stress Scale (NSS)

The NSS was developed to evaluate stress levels in preterm infants through behavioral and physiological indicators [23]. The scale consists of eight subscales: facial expression, skin color, respiration, activity, comfort, muscle tone, extremities, and posture. Each item is scored on a 3-point scale ranging from 0 to 2, where higher scores indicate greater stress-related responses. The total score ranges from 0 to 16, with higher total scores reflecting higher levels of neonatal stress. During the assessment, infants are observed for stress-related behavioral and physiological signs, and scores are assigned according

to the criteria defined for each subscale. The NSS has been used as a practical tool for evaluating stress responses in preterm infants in clinical settings.

Actigraphy (SOMNOWatch™)

An actigraphy device was used to assess the sleep process. Actigraphy is a non-invasive method that continuously records limb movement and quantitatively measures activity levels at defined time intervals. Data obtained from the device were analyzed in 1-minute epochs using the manufacturer-provided analysis software, with sleep and wakefulness states automatically classified. Sleep parameters were calculated as follows: sleep efficiency, total sleep duration, number of awakenings, and wake after sleep onset (WASO). Actigraphy has been reported to be a widely used and validated method for assessing sleep-wake patterns in infant and young child populations, with its validity supported in comparative studies with polysomnography [24].

Procedure and Data Collection

Prior to study commencement, parents were informed about the purpose of the study and written informed consent was obtained. Participants were assigned to the intervention and control groups via randomization. Data were collected at baseline and at the conclusion of the three-day intervention period using identical protocols.

Intervention Group: Massage Protocol

Massage was administered by a certified researcher with specialized training in infant massage over three consecutive days, three times daily (08:00–10:00, 14:00–16:00, and 19:00–21:00), either one hour before or one hour after feeding.

Each session lasted 15 minutes and was structured to incorporate both tactile (effleurage, petrissage) and kinesthetic (passive joint movements) stimulation (5 min tactile + 5 min kinesthetic + 5 min tactile). Infants were positioned in the supine and prone positions sequentially during the massage.

Prior to each session, hand hygiene was performed, hands were warmed, and the incubator temperature was maintained between 30°C and 36°C. Environmental stimuli (light and noise) were minimized throughout the session to preserve infant comfort.

Infants in the intervention group underwent 24-hour actigraphy recordings before the intervention and after the third day, with sleep duration, sleep efficiency, number of awakenings, and WASO parameters assessed. Vital signs were recorded and scales administered at the same time points.

Control Group: Standard Care

Infants in the control group received the standard nursing care protocol of the NICU. This care encompassed feeding, positioning, vital sign monitoring, hygiene maintenance, and required medical treatments; no additional tactile stimulation or massage was provided during the study period. Parallel to the intervention group, 24-hour actigraphy recordings were conducted for the control group at baseline and at the end of the third day; vital signs were recorded and scale assessments were performed at the same time points.

Ethical Considerations

Before initiation of the study, approval was obtained from the Non-Interventional Clinical Research Ethics Committee of Yüzüncü Yıl University (Approval No: 2024/10-14, Date: 20.09.2024) and institutional

authorization was secured from the hospital administration. Parents of the infants were provided with a written informed consent form and both written and verbal consent were obtained. The study was conducted in adherence to the principles of the Declaration of Helsinki throughout its entire duration.

Statistical Analysis

Data were analyzed using SPSS v27.0. Descriptive statistics included counts, percentages, means, and standard deviations. Normality of the data distribution was assessed via Skewness-Kurtosis values and the Shapiro-Wilk test. Between-group homogeneity was evaluated using the Chi-square and Fisher's exact tests; within-group changes were examined with the Paired Samples t-test, and between-group differences with the Independent Samples t-test. A significance level of $P < 0.05$ was applied to all analyses.

RESULTS

The baseline characteristics of the enrolled mothers and neonates are presented in Table 1. Statistical analyses revealed no significant differences between

TABLE 1. Baseline Characteristics of Mothers and Neonates

		Study group		Test statistics	P-value
		Massage	Control		
Maternal age (years)		30±6.87	28.15±6.15	0.898	0.375*
History of preterm birth (yes)	Yes	3 (15)	4 (20)	0.173	0.677**
	No	17 (85)	16 (80)		
Gestational age (weeks)	35.	9 (45)	10 (50)	0.497	0.780**
	36.	7 (35)	5 (25)		
	37.	4 (20)	5 (25)		
Birth weight (g)		2390.25±449.10	2390.25±449.10	1.171	0.249*
Infant sex	Girl	7 (35)	12 (60)	2.506	0.113**
	Boy	13 (65)	8 (40)		
Mode of delivery	Normal	3 (15)	3 (15)	0.000	1.000**
	C/S	17 (85)	17 (85)		
Feeding method	Breastfeeding	14 (70)	8 (40)	3.636	0.057**
	OG	6 (30)	12 (60)		

Data are shown as mean±standard deviation or n (%) where appropriate. C/S, caesarean section; OG, Orogastric tube.

*Independent Samples t-test, **Chi-square test.

the massage group (n=20) and the control group (n=20) with respect to maternal age, history of preterm birth, gestational age, birth weight, infant sex, mode of delivery, or feeding method (P>0.05). These findings confirm that the groups were homogeneously distributed across key clinical and demographic variables, thereby establishing an appropriate baseline for a randomized controlled trial.

No statistically significant difference in heart rate was detected between the massage and control groups at baseline (P=0.480). Following the intervention, a highly significant reduction in heart rate was observed in the massage group (from 143.20±10.55 to 128.15±7.91 bpm; P<0.001), whereas the reduction in the control group was comparatively modest (P=0.037). Post-intervention between-group comparisons demonstrated that the massage group had significantly lower heart rate values than the control group (t=3.463; P=0.001), with a large effect size (Cohen's d >0.8) (Table 2).

At baseline, SpO₂ values were significantly higher in the control group than in the massage group

(P=0.033). Following the intervention, a highly significant increase in oxygen saturation was recorded in the massage group (from 94.60±1.76% to 96.95±0.83%; P<0.001), whereas no significant change was observed in the control group (P=0.154). In the post-intervention period, SpO₂ levels in the massage group were statistically significantly higher than those in the control group (P<0.001, Table 2).

A significant reduction in respiratory rate was observed in the massage group post-intervention (P<0.001), whereas no significant change was detected in the control group (P=0.428). Between-group comparisons confirmed that the post-intervention respiratory rate was significantly lower in the massage group than in the control group (P=0.001). No statistically significant differences were found in body temperature, either between groups (P=0.567) or within groups over time (P>0.05) (Table 2).

Comparative analysis of PICS total scores before and after the intervention is presented in Table 3. At baseline, no significant difference in comfort scores was found between the massage and control groups

TABLE 2. Comparison of Physiological Parameters by Group

	Massage	Control	Test statistic	P-value*
Pre-procedure peak heart rate	143.20±10.55	140.80±10.75	0.713	0.480
Post-procedure peak heart rate	128.15±7.91	137.65±9.38	3.463	0.001
Test statistic	8.965	2.244		
P-value**	<0.001	0.037		
Pre-procedure oxygen saturation	94.60±1.76	95.85±1.81	2.212	0.033
Post-procedure oxygen saturation	96.95±0.83	95.35±1.63	3.914	<0.001
Test statistic	7.378	1.486		
P-value**	<0.001	0.154		
Pre-procedure respiratory rate	53.90±1.21	53.00±1.38	2.196	0.034
Post-procedure respiratory rate	52.00±0.65	53.20±1.36	3.559	0.001
Test statistic	7.025	0.809		
P-value**	<0.001	0.428		
Pre-procedure fever	36.89±0.13	36.95±0.08	1.696	0.098
Post-procedure temperature	36.87±0.15	36.90±0.13	0.578	0.567
Test statistic	0.476	1.278		
P-value**	0.640	0.217		

Data are shown as mean±standard deviation.

*Independent Samples t-Test; **Paired Samples t-Test. Statistically significant P-values are shown in bold.

TABLE 3. Comparison of the Premature Infant Comfort Scale by Group

	Study status		Test statistics	P-value*
	Massage	Control		
Pre-procedure comfort total score	14.75±1.52	14.4±1.47	0.742	0.460
Post-procedure comfort total score	10.65±2.52	13.15±2.28	3.292	0.002
Test statistic	9.853	2.544		
P-value**	<0.001	0.020		

Data are shown as mean±standard deviation.

*Independent Samples t-test; **Paired Samples t-test. Statistically significant P-values are shown in bold.

(14.75±1.5 and 14.4±1.47, respectively, $P=0.460$), confirming group homogeneity at study onset. Following the intervention, comfort scores in the massage group were significantly lower (indicating higher comfort) than in the control group (10.65±2.52 vs. 13.15±2.28; $t=3.292$; $P=0.002$). A highly significant within-group reduction (improvement) in comfort scores was recorded in the massage group post-intervention ($P<0.001$). Although the control group also demonstrated a statistically significant reduction in scores over time ($P=0.020$), the clinical depth and statistical power of this improvement were considerably less pronounced than in the massage group.

Neonatal Stress Scale scores were analyzed both between and within groups (Table 4). At baseline, no significant difference was found between the massage group (4.15±2.01) and the control group (4.35±2.08) ($P=0.759$), indicating comparable stress levels prior to the intervention period. Following the study period, the mean NSS score decreased to 2.00±1.62 in the massage group and to 2.95±2.04 in the control group. Although the massage group demonstrated lower post-procedure stress scores, the between-group difference

was not statistically significant ($t=1.631$, $P=0.111$). Within-group analyses showed a significant reduction in NSS scores from pre-procedure to post-procedure measurements in both groups (massage group: $P<0.001$; control group: $P=0.005$).

Findings pertaining to sleep efficiency, total sleep duration, number of awakenings, and WASO are presented in Table 5. Within-group analyses revealed statistically significant improvements in all sleep parameters post-intervention relative to baseline in both the massage and control groups ($P<0.001$ for most parameters). Sleep efficiency in the massage group increased from 83.25% to 88.5%, while a comparable improvement was observed in the control group (from 84.6% to 87.9%).

Between-group comparisons in the post-intervention period revealed no statistically significant differences in sleep efficiency ($P=0.776$), sleep duration ($P=0.308$), number of awakenings ($P=0.656$), or WASO ($P=0.585$). These findings indicate that both interventions produced positive effects on sleep parameters; however, massage did not demonstrate superior efficacy compared to the control condition.

TABLE 4. Comparison of Neonatal Stress Scale Scores by Group

	Study group		Test statistics	P-value*
	Massage	Control		
Pre-procedure NSS total	4.15±2.01	4.35±2.08	0.309	0.759
Post-procedure NSS total	2.00±1.62	2.95±2.04	1.631	0.111
Test statistic	4.454	3.200		
P-value**	0.001	0.005		

Data are shown as mean±standard deviation. NSS, neonatal stress scale.

*Independent Samples t-test; **Paired Samples t-test. Statistically significant P-values are shown in bold.

TABLE 5. Comparison of Sleep Parameters in Infants by Group

	Study group		Test statistics	P-value*
	Massage	Control		
Pre-procedure sleep efficiency	83.25±10.42	84.6±9.07	0.437	0.665
Post-procedure sleep efficiency	88.5±7.02	87.9±6.17	0.287	0.776
Test statistic	4.794	3.334		
P-value**	<0.001	0.003		
Pre-procedure sleep duration	837.8±179.87	856.85±203.49	0.314	0.755
Post-procedure sleep duration	968.8±145.94	918.35±162.2	1.034	0.308
Test statistic	6.550	2.707		
P-value**	<0.001	0.014		
Number of awakenings before surgery	97.75±58.45	95.2±66.18	0.129	0.898
Post-procedure awakening count	72.8±48.69	80.7±61.8	0.449	0.656
Test statistic	6.814	5.139		
P-value**	<0.001	<0.001		
Pre-procedure WASO value	121.45±71.96	118.8±83.03	0.108	0.915
Post-procedure WASO value	84.95±58.82	96.6±74.1	0.551	0.585
Test statistic	7.541	5.910		
P-value**	<0.001	<0.001		

Data are shown as mean±standard deviation. WASO, wake after sleep onset.

*Independent Samples t-test; **Paired Samples t-test. Statistically significant P-values are shown in bold.

DISCUSSION

This randomized controlled trial comprehensively assessed the effects of structured massage therapy on physiological stabilization, comfort, stress, and sleep parameters in late preterm infants (35–37 gestational weeks). The findings demonstrate that massage therapy produced statistically and clinically significant improvements in heart rate, oxygen saturation, and respiratory rate. Comfort levels were significantly higher in the massage group than in the control group; within-group improvements were observed in stress and sleep parameters in both groups, although no statistically significant between-group differences were identified. This pattern is largely consistent with the existing literature and yields important insights into the distinctive clinical characteristics of the late preterm population.

In the present study, massaged infants demonstrated a significant reduction in heart rate from 143.20±10.55 to 128.15±7.91 bpm ($P<0.001$), a

significant increase in oxygen saturation from 94.60% to 96.95% ($P<0.001$), and a marked decrease in respiratory rate ($P<0.001$). Changes in these parameters were considerably more limited in the control group, with post-intervention between-group differences reaching statistical significance ($P=0.001$ for heart rate; $P<0.001$ for SpO_2). These findings directly corroborate those of the randomized controlled trial by Düken *et al.* [19], which applied the same three-day protocol and reported significant reductions in heart rate and respiratory rate, along with significant increases in oxygen saturation, in the massage group compared to controls. Concurrently, our physiological findings align with the randomized controlled trial by Alinejad-Naeini *et al.* [25], which demonstrated that the 'M' Technique[®] significantly improved physiological stabilization in preterm neonates — an effect the authors attributed to enhanced vagal activation within the polyvagal theory framework. A meta-analysis by Zhang *et al.* [26] encompassing 15 randomized controlled trials

confirmed that massage therapy significantly increased oxygen saturation (SMD=2.00; 95% CI: 1.17–2.83; $P<0.0001$). Furthermore, Weerakul *et al.* [27] highlighted that moderate-pressure massage suppresses sympathetic nervous system activity, activates vagal pathways, and thereby reduces heart rate while augmenting gastrointestinal motility in preterm infants.

The primary mechanism underlying these effects of massage on physiological stabilization is explained by parasympathetic system activation triggered by stimulation of cutaneous mechanoreceptors. Tactile stimulation activates the posterior insular cortex via C-tactile afferents, suppressing autonomic arousal and producing reductions in heart rate and respiratory rate [19]. The fact that our intervention protocol encompassed both tactile (effleurage, petrissage) and kinesthetic (passive joint movement) components supports the synergistic effect of this multimodal stimulation. In this regard, the systematic review and meta-analysis by Mollà-Casanova *et al.* [28] demonstrated that massage therapy combined with passive mobilizations significantly improved weight gain and length of hospitalization, further substantiating the clinical value of the kinesthetic component in our protocol. The stable body temperatures observed in both groups indicate that massage did not exert any adverse effect on thermoregulation, which is consistent with the safety profiles reported by Alinejad-Naeini *et al.* [25] and Zhang *et al.* [26].

The significant reduction in PICS total scores in the massage group (from 14.75 ± 1.52 to 10.65 ± 2.52 ; $P<0.001$) and the statistically significant post-intervention between-group difference ($P=0.002$) supported Hypothesis H2 and demonstrated that massage exerts a clinically important effect on comfort in late preterm infants. This finding is consistent with the randomized controlled trial by Zhang and Wang [29], which showed that mother-administered massage significantly improved growth parameters in infants born at 32–34 weeks and in which mothers also qualitatively reported that their infants appeared relaxed and displayed a tendency toward sleep during the massage. The concurrent occurrence of comfort improvement alongside physiological stabilization is clinically significant, as the heart rate and respiration/crying subscales of the PICS are directly

related to the physiological parameters that improved in parallel. Massage is known to trigger the release of oxytocin and endorphins through tactile stimulation; these neuropeptides attenuate pain perception and suppress autonomic arousal, thereby enhancing comfort [27]. The statistically significant improvement in comfort observed over time in the control group as well ($P=0.020$) suggests that routine NICU care provides a baseline level of comfort, although the improvement in the massage group was markedly superior in both effect size and clinical depth. This finding is further corroborated by the comprehensive umbrella review by Chen *et al.* [30] evaluating the effects of pediatric massage in infants and children under 5 years of age, which demonstrated significant effects of massage on pain management and oxygen saturation with high-quality evidence.

NSS assessments revealed significant reductions in stress scores post-intervention relative to baseline in both groups (massage group: $P<0.001$; control group: $P=0.005$); however, the between-group difference did not reach statistical significance post-intervention ($P=0.111$). While this finding does not support Hypothesis H3, the substantially greater statistical power of the stress reduction in the massage group ($P<0.001$) compared to the control group ($P=0.005$) suggests that a between-group difference may reach significance in a study with a larger sample size. This pattern is consistent with the discussion presented in the PREMM protocol study by Lai *et al.* [31], which highlighted the prevailing environmental stress burden in preterm infants and the spontaneous improvement afforded by routine NICU care.

Biochemical evidence regarding the effect of massage on stress further supports these findings. In the first systematic review and meta-analysis to evaluate the effect of massage on cortisol levels in infants, Jabraeili *et al.* [32] reported mixed cortisol responses, yet noted that some studies demonstrated significant reductions in cortisol levels following massage. This meta-analysis highlighted methodological inconsistencies with respect to biochemical markers and underscored the need for studies incorporating standardized cortisol measurement. McCarty *et al.* [33] measured salivary cortisol levels in extremely preterm infants and their mothers before and after massage, reporting a significant cortisol reduction of 20.1% in responsive

mothers — a proportion that exceeded the 15.5% threshold used to define a clinically meaningful stress response. The authors noted that parent-administered massage may reduce the physiological stress biomarker in mothers in the short term, highlighting important implications for its integration into NICU clinical practice. A key methodological limitation of the present study is that stress assessment relied solely on observational NSS scores, without the inclusion of biochemical biomarkers such as cortisol or salivary alpha-amylase, which constrains more objective interpretation of the stress response. Future studies incorporating such biomarkers alongside the NSS may more definitively elucidate the specific effects of massage on stress mechanisms.

Significant improvements were recorded in all sleep parameters — sleep efficiency, total sleep duration, number of awakenings, and WASO — relative to baseline in both groups ($P < 0.001$); however, post-intervention between-group comparisons did not yield statistical significance ($P > 0.05$). This finding does not support Hypothesis H4; nonetheless, the increase in total sleep duration in the massage group from 837.8 ± 179.87 to 968.8 ± 145.94 minutes and the improvement in sleep efficiency from 83.25% to 88.5% should be regarded as clinically meaningful changes. The study by Düken and Yayan [17] provides clear context for these findings: in preterm infants aged 28–37 weeks, sleep duration in the massage group increased by approximately five hours compared to the pre-intervention period, whereas sleep duration decreased in the control group ($P < 0.001$). In that study, although the between-group difference in sleep duration was significant, no significant differences were found in sleep efficiency, number of awakenings, or WASO, suggesting that individual sleep parameters are sensitive to distinct NICU environmental factors and intervention duration.

Multiple explanations exist for the failure to achieve between-group significance for sleep in the present study. First, the more mature nervous systems of late preterm infants (35–37 weeks) facilitate spontaneous sleep improvement in response to standard NICU care, which accounts for the strong within-group improvements also observed in the control group. Additionally, the three-day intervention period may have been insufficient to achieve between-

group statistical significance for sleep parameters; notably, Lai *et al.* [31] demonstrated that long-term massage sustained until term-equivalent age positively influenced cerebral electrical activity in very preterm infants. Moreover, the sample size of $n=20$ per group in the present study may not have provided adequate statistical power for sleep variables, despite the objective measurement afforded by SOMNOwatch™ actigraphy. The partial validation of actigraphy against polysomnography in the preterm infant population must also be considered when interpreting results [24]. Recent literature supports these findings: Belpinar *et al.* [34] evaluated the effects of massage and foot reflexology on sleep parameters in preterm infants aged 30–37 weeks using actigraphy and demonstrated that sleep efficiency and total sleep duration improved significantly in the massage group compared to the control group. Similarly, the meta-analysis by Bilgiç *et al.* [35] examining the effects of non-pharmacological interventions on sleep parameters in preterm infants emphasized that intervention duration and sample size play critical roles in determining between-group significance.

Strengths and Limitations

This study offers several methodological strengths that support the validity and clinical relevance of its findings. The randomized controlled trial design, implemented in strict accordance with CONSORT guidelines, minimized selection bias and ensured homogeneous group distribution across all key demographic and clinical variables. A distinguishing feature of this study is its simultaneous assessment of four distinct outcome domains — physiological stability, comfort, stress, and sleep — within a single protocol, enabling a comprehensive evaluation of massage therapy's multidimensional effects, an approach rarely adopted in the existing literature. The use of SOMNOwatch™ actigraphy for sleep measurement provided continuous, objective, and quantitative data across 24-hour periods, substantially reducing observer bias compared to conventional observational methods. Intervention fidelity was ensured through a standardized massage protocol administered by a certified researcher, enhancing reproducibility and translational applicability. Importantly, by specifically targeting the late preterm

cohort (35–37 gestational weeks) — a clinically underrepresented yet high-risk population in neonatal massage research — this study contributes population-specific evidence to the field. Finally, assessor blinding during statistical analysis mitigated measurement bias, preserving analytical integrity within the inherent constraints of this intervention type.

The single-center design limits the generalizability of findings to different clinical and cultural settings. Additionally, the relatively small sample size further limits the statistical power and generalizability of the findings. Blinding of the interventionist was not achievable given the nature of the intervention. The restriction of stress assessment to a behavioral observation scale — without biomarkers such as cortisol, which could not be utilized due to limitations in hospital infrastructure — represents an additional limitation. Furthermore, the sensitivity of biomarker levels in preterm infants to physiological and environmental factors may complicate standardization of measurements. These findings are generalizable to clinically stable, late preterm infants at 35–37 gestational weeks cared for in settings comparable to the study institution. Generalizability is limited for infants at lower gestational ages, clinically unstable cases, and centers employing different care protocols.

CONCLUSION

The findings of this study provide clinically relevant evidence for NICU protocols that prioritize physiological stabilization and comfort optimization. Structured massage therapy resulted in significant improvements in key physiological parameters and infant comfort, supporting its role as an effective, evidence-based, non-pharmacological intervention in late preterm infants. While improvements in sleep and stress parameters were observed within groups, the absence of significant between-group differences suggests that these outcomes may be influenced by routine NICU care and the relatively short intervention duration. Therefore, the primary clinical value of massage therapy in this population appears to lie in its capacity to enhance physiological stability and comfort rather than to produce additional benefits in sleep and stress beyond standard care. Integration of massage therapy into routine neonatal

nursing care is important for improving physiological parameters. The inclusion of parents in the massage process may not only contribute to improvements in the infant's clinical trajectory but may also strengthen mother-infant bonding and parental self-efficacy. In future studies, larger sample sizes, longer intervention durations (at least 7–14 days), multicenter designs, and the addition of cortisol measurements and EEG-based sleep assessments could more definitively elucidate the mechanisms of action and stress-sleep outcomes of massage therapy. Gestational age-specific designs comparing late preterm infants with those born at earlier gestational ages are also expected to contribute to identifying the developmental threshold below which the biological and clinical effects of the intervention become most pronounced.

Ethics Approval and Consent to Participate

This study was approved by the Van Yüzüncü Yıl University Non-interventional Clinical Research Ethics Committee (Decision No: 2024/10-4; date: 20.09.2024). All procedures were conducted in accordance with the ethical standards of the institutional and national research committee and with the 1964 Helsinki Declaration and its later amendments. Written informed consent was obtained from all participants' parents for the publication of this study.

Trial Registration

This randomized controlled trial was registered in clinicaltrials.gov, and the registration number NCT07322731 was received.

Data Availability

All data generated or analyzed during this study are included in this published article. The data that support the findings of this study are available on request from the corresponding author, upon reasonable request.

Authors' Contribution

Study Conception: RA, YG, MED; Study Design: RA, YG, MED; Supervision: RA; Funding: YYU BAP (Project No: 11429); Materials: RA, YG, MED; Data Collection and/or Processing: RA, YG; Statistical Analysis and/or Data Interpretation: RA, MED; Literature Review: RA, YG, MED; Manuscript Preparation: RA, YG, MED; and Critical Review: RA.

Conflict of Interest

The author(s) disclosed no conflict of interest during the preparation or publication of this manuscript.

Funding

This study was supported by the Scientific Research Projects Unit (BAP) of Van Yüzüncü Yıl University under a Career Start Project (Project No: 11429).

Acknowledgments

The authors would like to express their sincere gratitude to all preterm infants' caregivers who participated in this study for their valuable time and contributions.

Generative Artificial Intelligence Statement

The authors used DeepL for initial translation and ChatGPT for language editing and proofreading to improve the clarity and readability of the manuscript. These generative AI tools were used solely for linguistic enhancement. All scientific content, including study design, data analysis, interpretation, and conclusions, was developed entirely by the authors. The authors take full responsibility for the integrity and accuracy of the work.

Editor's Note

All statements made in this article are solely those of the authors and do not represent the views of their affiliates or the publisher, editors, or reviewers. Any claims made by any product or manufacturer that may be evaluated in this article are not guaranteed or endorsed by the publisher.

REFERENCES

- World Health Organization. Preterm birth and low birth weight [Internet]. Geneva: World Health Organization. Accessed date: March 1, 2026. Available from: <https://www.who.int/teams/maternal-newborn-child-adolescent-health-and-ageing/newborn-health/preterm-and-low-birth-weight>
- Çelik F. Preterm Bebeklerde Oral Stimulasyon ve Emzirme Destek Sisteminin Tam Anne Memesine Geçiş Süresi ve Emme Başarısı Üzerine Etkisi: Randomize Kontrollü Çalışma [Effects of Oral Stimulstion and Breastfeeding Support System on the Transition to Time to Full Breast of Mother and Sucking Success in Preterm Infants: A Randomized Controlled Trial] [master's thesis]. Celal Bayar Üniversitesi Sağlık Bilimleri Enstitüsü, Manisa, 2019. [Article in Turkish]
- Turan T, Erdoğan Ç. Yenidoğan Yoğun Bakım Ünitesindeki Prematüre Bebeğin Gelişiminin Desteklenmesi [Supporting the Development of Premature Babies in Neonatal Intensive Care Unit]. GOP Taksim EAH JAREN. 2018;4(2):127-132. doi: 10.5222/jaren.2018.127. [Article in Turkish]
- Tekgündüz KŞ, Polat S, Gürol A, Apay SE. Oral Glucose and Listening to Lullaby to Decrease Pain in Preterm Infants Supported with NCPAP: A Randomized Controlled Trial. Pain Manag Nurs. 2019;20(1):54-61. doi: 10.1016/j.pmn.2018.04.008.
- Cardoso SM, Kozłowski Lde C, Lacerda AB, Marques JM, Ribas A. Newborn physiological responses to noise in the neonatal unit. Braz J Otorhinolaryngol. 2015;81(6):583-588. doi: 10.1016/j.bjorl.2014.11.008.
- da Silva CM, Caçao JM, Silva KC, Marques CF, Merey LS. Physiological responses of preterm newborn infants submitted to classical music therapy. Rev Paul Pediatr. 2013;31(1):30-36. doi: 10.1590/s0103-05822013000100006.
- Park J. Sleep Promotion for Preterm Infants in the NICU. Nurs Womens Health. 2020;24(1):24-35. doi: 10.1016/j.nwh.2019.11.004.
- Yates CC, Mitchell AJ, Booth MY, Williams DK, Lowe LM, Whit Hall R. The effects of massage therapy to induce sleep in infants born preterm. Pediatr Phys Ther. 2014;26(4):405-410. doi: 10.1097/PEP.0000000000000081.
- Güler E, Aydın Ateş N, Kügçümen G. Kanguru Bakımının Düşük Doğum Ağırlıklı ve Prematüre Yenidoğana Etkileri [Effects of Kangaroo Care on Low Birth Weight and Premature Newborn]. Cumhuriyet Üniv Sağ Bil Enst Derg. 2019;4(2):1-5. [Article in Turkish]
- Kanbur BN, Balcı S. Preterm Yenidoğanlarda Koku [Olfaction in Preterm Newborn]. Sağlık Bilim Meslekleri Derg. 2017;4(3):272-276. doi: 10.17681/hsp.277840.
- Field T. Massage therapy research review. Complement Ther Clin Pract. 2016;24:19-31. doi: 10.1016/j.ctcp.2016.04.005.
- Firmino C, Rodrigues M, Franco S, et al. Nursing Interventions That Promote Sleep in Preterm Newborns in the Neonatal Intensive Care Units: An Integrative Review. Int J Environ Res Public Health. 2022;19(17):10953. doi: 10.3390/ijerph191710953.
- de Britto Pereira PAD, Mendes Abdala CV, Portella CF, Ghelman R, Schweitzer MC. Pediatrics massage evidence map. Complement Ther Med. 2021;61:102774. doi: 10.1016/j.ctim.2021.102774.
- Akhavan Karbasi S, Golestan M, Fallah R, Golshan M, Dehghan Z. Effect of body massage on increase of low birth weight neonates growth parameters: A randomized clinical trial. Iran J Reprod Med. 2013;11(7):583-588.
- Gonzalez AP, Vasquez-Mendoza G, García-Vela A, Guzmán-Ramirez A, Salazar-Torres M, Romero-Gutierrez G. Weight gain in preterm infants following parent-administered Vimala massage: a randomized controlled trial. Am J Perinatol. 2009;26(4):247-252. doi: 10.1055/s-0028-1103151.
- Doğan P, Akçay N, Bal Yılmaz H. Pediatri Hemşireliğinde Önemli Bir Girişim: Atravmatik Bakım ve Yaklaşımları [An Important Initiative in Pediatric Nursing: Atraumatic Care and Approaches]. İzmir Kâtip Çelebi Univ Sağlık Bilim Fak Derg. 2021;6(2):131-135. [Article in Turkish]
- Düken ME, Yayan EH. The effects of massage therapy and

- white noise application on premature infants' sleep. *Explore (NY)*. 2024;20(3):319-327. doi: 10.1016/j.explore.2023.09.002.
18. Açar SÇ, Bal Yılmaz H. Preterm Bebeklerde Masajın Vital Bulgular ve Büyüme Parametrelerine Etkisi: Yarı Deneysel Çalışma [Effects of Massage on Vital Signs and Growth Parameters in Preterm Infants: The Quasi-Experimental Study]. *Türkiye Klinikleri J Nurs Sci*. 2022;14(3):648-658. doi: 10.5336/nurses.2021-86377.
19. Düken ME, Yayan EH, Arslan Z. The effects of massage and white noise on physiological parameters of preterm infants: A randomized controlled research. *J Neonatal Nurs*. 2024;30(4):360-367. doi: 10.1016/j.jnn.2023.11.008
20. Ambuel B, Hamlett KW, Marx CM, Blumer JL. Assessing distress in pediatric intensive care environments: the COMFORT scale. *J Pediatr Psychol*. 1992;17(1):95-109. doi: 10.1093/jpepsy/17.1.95.
21. Caljouw MAA, Kloos MAC, Olivier MY, et al. Measurement of pain in premature infants with a gestational age between 28 to 37 weeks: Validation of the adapted COMFORT scale. *J Neonatal Nurs*. 2007;13(1):13-18. doi: 10.1016/j.jnn.2006.11.007.
22. Küçük Alemdar D, Güdücü Tüfekçi F. Prematüre Bebek Konfor Ölçeği'nin Türkçe Geçerlilik ve Güvenilirliği [The Reliability and Validity of the Premature Infant Comfort Scale's Turkish]. *Hemşirelik Eğitim ve Araştırma Dergisi*. 2015;12(2):142-148. doi: 10.5222/HEAD.2015.142. [Article in Turkish]
23. Ceylan SS, Bolışık B. Yenidoğan Stres Ölçeği'nin Psikometrik Özelliklerinin İncelenmesi [Examining Psychometric Properties of Newborn Stress Scale]. *ACU Sağlık Bil Derg* 2017(2):97-103. [Article in Turkish]
24. Schoch SF, Kurth S, Werner H. Actigraphy in sleep research with infants and young children: Current practices and future benefits of standardized reporting. *J Sleep Res*. 2021;30(3):e13134. doi: 10.1111/jsr.13134.
25. Alinejad-Naeini M, Heidari-Beni F, Mohagheghi P, Sohrabi S, Jeshvaghani SS. The effect of the 'M' Technique® massage on physiological parameters in preterm neonates: a randomized controlled trial study. *J Pediatr Neonatal Individ Med*. 2022;11(2):e110212. doi: 10.7363/110212.
26. Zhang Y, Duan C, Cheng L, Li H. Effects of massage therapy on preterm infants and their mothers: a systematic review and meta-analysis of randomized controlled trials. *Front Pediatr*. 2023;11:1198730. doi: 10.3389/fped.2023.1198730.
27. Weerakul J, Apiraknapanon Y, Sanjaiban M, Intasen S, Tipsuwan S, Bhumipraphat M. The Effects of Infant Massage Therapy on Preterm Neonatal Outcomes: A Clinical Trial. *Int J Pediatr*. 2025;2025:2451284. doi: 10.1155/ijpe/2451284.
28. Mollà-Casanova S, Sempere-Rubio N, Muñoz-Gómez E, Aguilar-Rodríguez M, Serra-Añó P, Inglés M. Effects of massage therapy alone or together with passive mobilisations on weight gain and length of hospitalisation in preterm infants: Systematic review and meta-analysis. *Early Hum Dev*. 2023;182:105790. doi: 10.1016/j.earlhumdev.2023.105790.
29. Zhang X, Wang J. Massage intervention for preterm infants by their mothers: A randomized controlled trial. *J Spec Pediatr Nurs*. 2019;24(2):e12238. doi: 10.1111/jspn.12238.
30. Chen SC, Lin SL, Wang M, et al. Pediatric massage therapy in infants and children under 5 years: An umbrella review of systematic reviews. *Heliyon*. 2024;10(16):e35993. doi: 10.1016/j.heliyon.2024.e35993.
31. Lai MM, D'Acunto G, Guzzetta A, et al. PREMM: preterm early massage by the mother: protocol of a randomised controlled trial of massage therapy in very preterm infants. *BMC Pediatr*. 2016;16(1):146. doi: 10.1186/s12887-016-0678-7.
32. Jabraeili M, Mirghafourvand M, Montazeri M, Farshi MR. The effect of massage on cortisol level in infants: A systematic review and meta-analysis. *J Neonatal Nurs*. 2023;29(5):720-727. doi: 10.1016/j.jnn.2023.02.010.
33. McCarty D, Silver R, Quinn L, Dusing S, O'Shea TM. Infant massage as a stress management technique for parents of hospitalized extremely preterm infants. *Infant Ment Health J*. 2024;45(1):11-21. doi: 10.1002/imhj.22095.
34. Belpınar A, Yayan EH, Özdemir A. The Effect of Massage and Foot Reflexology on Sleep in Newborns. *Biol Res Nurs*. 2025;27(3):504-517. doi: 10.1177/10998004251332911.
35. Bilgiç FŞ, Karaahmet AY, Okay B. Effects of non-pharmacological interventions on sleep in premature infants: meta-analysis. *Rev Assoc Med Bras (1992)*. 2024;70(11):e20241187. doi: 10.1590/1806-9282.20241187.
36. Grant S, Mayo-Wilson E, Montgomery P, et al. on behalf of the CONSORT-SPI Group. CONSORT-SPI 2018 Explanation and Elaboration: guidance for reporting social and psychological intervention trials. *Trials*. 2018;19(1):406. doi: 10.1186/s13063-018-2735-z.